
AD-R178 1	L67 I	THE PLASP SIMPLE AF BUFFALO D JBUFFALO	ION DIS PROXIN EPT OF DC/87/	PERSI INTIO. CHEM TR-31	ON REL . (U) S ISTRY NOOC!	ATION TATE D A 4-86-	ON A UMIY O JELSKI K- 00 43	ROUGH F MEN ET AL	SURFA YORK MAR F/G	CE: R AT 87 20/9	S/	1
											1 N () 	
		_										



OFFICE OF NAVAL RESEARCH

Contract N00014-K-0043

R & T Code 413f0001---01

TECHNICAL REPORT No. 31

The Plasmon Dispersion Relation on a Rough Surface: A Simple Approximation

by

Daniel Jelski and Thomas F. George

Prepared for Publication

in

Journal of Physical Chemistry

Departments of Chemistry and Physics State University of New York at Buffalo Buffalo, New York 14260

March 1987

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.



REPORT DOCUMENTATION PAGE							
1a. REPORT SECURITY CLASSIFICATION Unclassified	1b. RESTRICTIVE MARKINGS						
28. SECURITY CLASSIFICATION AUTHORITY	3. DISTRIBUTION/AVAILABILITY OF REPORT						
26. DECLASSIFICATION/DOWNGRADING SCHED	Approved for public release; distribution unlimited						
4. PERFORMING ORGANIZATION REPORT NUM	BER(S)	5. MONITORING OF	IGANIZATION RI	EPORT NUMB	ER(S)		
UBUFFALO/DC/87/TR-	31						
Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (If applicable)	7s. NAME OF MONITORING ORGANIZATION					
6c. ADDRESS (City, State and ZIP Code)		7b. ADDRESS (City, State and ZIP Code)					
Fronczak Hall, Amherst Campus Buffalo, New York 14260		Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217					
BA. NAME OF FUNDING/SPONSORING	85. OFFICE SYMBOL (if applicable)	9. PROCUREMENT			N NUMBER		
Office of Naval Research	, 4,2	Contract N00014-86-K-0043					
Bc. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUI	NDING NOS.	T			
Chemistry Program 800 N. Quincy Street		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT		
Arlington, Virginia 22217				<u> </u>			
The Plasmon Dispersion Relation on a Rough Surface: A Simple Approximation							
12 PERSONAL AUTHOR(S) Daniel A. Je	lski and <u>Thomas</u>	F. George					
13a. TYPE OF REPORT 13b. TIME C	OVERED	14. DATE OF REPOR		16. PA	GE COUNT 18		
16. SUPPLEMENTARY NOTATION							
Prepared for Publication	on in the Journa	l of Physical	Chemistry				
17. COSATI CODES	18 SUBJECT TERMS (C.) PLASMON DISPER						
FIELD GROUP SUB. GR.	ROUGH SURFACE	, P≠POLARIZED LIGHT;					
	CHEMICAL VAPOR		COMP	LEX DIELE	CTRIC CONSTANT		
This paper is concerned with periodic, laser-induced, chemical vapor deposition recently observed experimentally. In order to inquire further into this phenomena, it is first necessary to find a simple means of calculating the plasmon field strength for relatively deep gratings. The Rayleigh hypothesis is assumed, and only p-polarized, normally incident light is considered. A closed-form equation for the plasmon field intensity is then derived. Also discussed is the behavior of the plasmon dispersion relation for a shallow grating, but for a complex dielectric constant where the imaginary part is not necessarily small.							
20. DISTRIBUTION/AVAILABILITY OF ABSTRA	21. ABSTRACT SECURITY CLASSIFICATION						
UNCLASSIFIED/UNLIMITED 🖾 SAME AS RPT.	Unclassifi	ed					
224 NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nels	225. TELEPHONE N (Include Area Co (202) 696-44	ode)	22c. OFFICE	SYMBOL			

gyesten berestenen beveration akonomisten kontraktion kantakte sakstationen bestektionen

Journal of Physical Chemistry, in press

THE PLASMON DISPERSION RELATION ON A ROUGH SURFACE: A SIMPLE APPROXIMATION

Daniel A. Jelski and Thomas F. George

Departments of Chemistry and of Physics & Astronomy 239 Fronczak Hall State University of New York at Buffalo Buffalo, New York 14260

ABSTRACT

This paper is concerned with periodic, laser-induced, chemical vapor deposition recently observed experimentally. In order to inquire further into this phenomena, it is first necessary to find a simple means of calculating the plasmon field strength for relatively deep gratings. The Rayleigh hypothesis is assumed, and only p-polarized, normally incident light is considered. A closed-form equation for the plasmon field intensity is then derived. Also discussed is the behavior of the plasmon dispersion relation for a shallow grating, but for a complex dielectric constant where the imaginary part is not necessarily small.

Accesio						
NTIS	CRA&I	Ø				
DTIC	TAB					
Unanno	ounced	IJ				
Justific						
By						
A-1						
	NTIS DTIC Unanno Justific By Distrib.	Distribution / Availability Avail a	NTIS CRA&I DID TIC TAB Unannounced Ustrication By Distribution Availability Codes			

DTIC Enpy INSPECTED

1. Introduction

and tradition properties the second seconds assessed assessed assessed assessed the seconds. The seconds are seconds assessed assessed the seconds as a second assessed the second assessed to second assessed the second assessed to the second assessed the second asset the second asset the second asset the second asset

Our laboratory has been investigating laser-induced chemical vapor deposition processes as recently observed experimentally. 1-3 In particular, we have been interested in the periodic structures observed by Brueck and Ehrlich³. In a later paper, we shall discuss this experiment directly. This paper, however, will be devoted to developing the theoretical framework necessary for the calculations.

The periodic structure that is observed evolves from the plasmon field induced by the laser. The laser interacts with the random surface roughness. Because the plasmon is resonant at one frequency (for shallow gratings), that frequency component grows as

$$\frac{\partial \xi}{\partial t} = aI(x), \tag{1}$$

where ξ is the grating amplitude, I(x) is the total field intensity as a function of position, and a is the proportionality constant. From this it is clear that we need to develop expressions for the plasmon field intensity as a function of grating height and wavenumber. We can simplify the problem considerably in all that follows by assuming perpendicular incidence, in which case the x-dependence of the intensity depends only on the plasmon field and not on the reflected or incident fields. We can simplify the problem even further by using the Rayleigh expansion. This is an exact solution above the selvedge region and an approximation within the selvedge region. However, we are primarily interested in field strengths above the selvedge region, so that this expansion is justified.

Many authors have discussed the interaction of radiation with a rough surface or grating. Petit⁴ has written a good introduction to the subject.

Maradudin⁵ has discussed the behavior of surface polaritons and plasmons (which together are what we are generically referring to as a surface plasmon). In particular, he has worked out the theory for randomly rough surfaces and has written expressions for the dispersion relation using both the Rayleigh expansion and the exact formula using Green's functions. We shall make much use of this work as we go along. Other authors have also made significant contributions. Jha, Kirtley and Tsang⁶ have developed the theory for a shallow grating. These authors have made extensive use of the work of Toigo, Marvin, Celli and Hill, ^{7,8} who have written the Rayleigh expansion in a remarkably simple form. Finally, Agassi and George ^{9,10} have developed a dressed Rayleigh expansion which eliminates many of the numerical difficulties encountered previously.

We are interested in the plasmon field intensity above the selvedge region for reasonably deep gratings. We need an expression as a function of both grating height and grating wavenumber. Since we shall also need the derivatives of this function, it is important to derive an expression in closed form. To date, all calculations for deep gratings have involved the inversion of large matrices. Given the conditions of our application, this is impractical. Therefore, we must develop an approximation which is still valid in the deep grating case, but which provides a closed form equation for the plasmon field strength. Two recent papers 11,12 have made progress in this field. In particular, the work by Weber 2 represents a generalization of our present result. We will compare our result with his in the conclusion.

We should also point out that there has been considerable work done on a closely related problem. Fauchet and Siegman 13 have discovered ripples as a result of laser annealing, and Sipe et al. 14 have developed an elegant

theory to account for this phenomena. But these are effects of surface damage rather than surface deposition, and hence the theoretical requirements are more stringent in that the Rayleigh hypothesis can no longer be applied.

But there is one additional problem. The experiment of Brueck and Ehrlich involved a plasmon wave along a cadmium surface. The dielectric constant for cadmium (at UV frequencies) is approximately

$$\varepsilon = -2.5 + 1.3i. \tag{2}$$

Most calculations, even for shallow gratings, have assumed that the imaginary part of the dielectric constant is either zero^{4,5}, or small.⁶ Since this is not the case for us, we shall digress a bit and discuss the effect on the dispersion relation as the imaginary part of the dielectric constant becomes large.

The remainder of this paper is organized as follows. In section 2 we shall discuss the dispersion relation as a function of dielectric constant. In section 3 we shall derive our approximate formula for the plasmon field strength. Section 4 contains a discussion of our results, including a numerical comparison of our approximation with the exact calculation.

2. <u>Dispersion Relation</u>

というとは

We can write the solution of the homogenous Helmholtz equation as

$$\underline{\underline{E}}_{\mathbf{g}} = \underline{E}_{\mathbf{i}} \underline{\underline{P}}_{\mathbf{f}} - \exp(\mathbf{i}[\mathbf{k}\mathbf{x} - \alpha_{\mathbf{0}}\mathbf{z}]) + \sum_{\mathbf{f} = -\infty}^{\infty} \mathbf{A}_{\mathbf{f}} \underline{\underline{P}}_{\mathbf{f}} + \exp(\mathbf{i}[\mathbf{k}_{\mathbf{f}}\mathbf{x} + \alpha_{\mathbf{f}}\mathbf{z}])$$

$$\mathbf{z} > \xi(\mathbf{x})$$
(3)

4

$$\underline{\underline{R}}_{b} = \sqrt{\varepsilon} \underline{s} \sum_{\underline{t}} C_{\underline{t}} \exp(i[k_{\underline{t}}x - \beta_{\underline{t}}z])$$

$$z < \xi(x), \qquad (4)$$

where

$$k = \frac{\omega}{c} \quad \epsilon = \epsilon_1 + i\epsilon_2 \qquad \epsilon_1 < -1$$

$$k_{\ell} = k_{t} + \ell k_{g} \qquad \alpha_{\ell} = (k^2 - k_{\ell}^2)^{1/2} \qquad \lim_{\epsilon \to 0} > 0 \qquad (5)$$

$$\beta_{\ell} = (\epsilon k^2 - k_{\ell}^2)^{1/2} \qquad \lim_{\epsilon \to 0} > 0 \qquad \underline{p}_{\ell \pm} = \frac{1}{k} [k_{\ell} \underline{z} + \alpha_{\ell} \underline{x}]$$

and where k_g is the plasmon frequency and E_i the incident field strength. The notation has been culled from refs. 5 and 9. This solution is exact as long as one is outside the selvedge region. Within the selvedge region it may be a good approximation, but as mentioned previously, we are primarily concerned with the region outside, and hence we can use eqs. 4 and 5 without reservation. From the above, it is apparent that A_0 is the reflected field, while A_1 is the 1-th order Bragg reflection (α_1 real), or surface plasmon (α_1 imaginary). We are interested in the plasmon effect, and hence we shall always assume that $k_g > k$.

Now, suppose that the surface profile is sinusoidal and can be written as

$$\xi(x) = \xi \cos(k_{g}x) , \qquad (6)$$

and that ξ is small. Then, for perpendicularly incident, p-polarized light, we can write the the plasmon field strength as

$$A_{\underline{t}} = \frac{\alpha_0 \beta_{\underline{t}} [k^2 \xi (\epsilon \alpha_{\underline{t}} - \beta_{\underline{t}})]}{\epsilon k^2 - k_{\underline{t}}^2 (1 + \epsilon)} . \qquad (7)$$

For small ξ , $\Lambda_{\underline{\xi}}$ is very small unless the denominator is very small. If ϵ is real, the denominator, which we shall call the resonance factor and denote by R, is zero if

$$k_g^2 = k^2 \frac{\epsilon_1}{\epsilon_1 + 1} . \tag{8}$$

This is the familiar plasmon resonance condition for a flat surface and a real dielectric constant. However, if ϵ is complex then the strength of the plasmon field depends on the modulus of R, and it is no longer sufficient merely to minimize the real part. We must then minimize

$$|R|^2 = [k_g^2(1+\epsilon_1) - k^2\epsilon_1]^2 + [\epsilon_2(k_g^2 - k^2)]^2.$$
 (9)

Differentiating with respect to k and setting the result to zero, we obtain a new dispersion relation

AND THE PROPERTY OF THE PROPER

$$k_g^2 = \frac{k^2[\epsilon_1(1+\epsilon_1) - \epsilon_2^2]}{(1+\epsilon_1)^2 + \epsilon_2^2}.$$
 (10)

As $\epsilon_2 \to 0$ this reduces to eq. 5. We also note that if $\epsilon_2 \neq 0$, then $|R| \neq 0$, and the resonance is both dampened and broadened. Equation 10 is the flat surface plasmon dispersion relation for an arbitrary, complex-valued dielectric constant.

As the grating becomes deeper, R also depends on higher powers of ξ_1 , whereby a branching occurs in the dispersion relation and the resonance becomes less pronounced. Thus the mechanism of the periodic deposition process should now be clear. In the beginning, the incident light is resonant with one frequency component of the randomly rough, but very shallow grating. Because of this resonance, the plasmon of that frequency is much stronger than the others. Hence the field intensity is periodic, and deposition, following eq. 1, has the same periodicity, leading to a growth in amplitude of the grating.

As the grating becomes larger, the dispersion relation branches, and the resonance is correspondingly broadened and dampened. This results in a decrease in the periodicity of the field intensity and a corresponding decrease in the differential rate of deposition (i.e., deposition occurs evenly, not necessarily more slowly). Hence the grating stops growing.

3. Calculation of Plasmon Field Intensity

We can write an expression for the coefficients of eq. 3 as 7,9

$$\sum_{\ell=-\infty}^{\infty} \mathbf{M}_{\mathbf{m}, \, \hat{\mathbf{z}}} \mathbf{A}_{\hat{\mathbf{z}}} = \boldsymbol{\mu}_{\mathbf{m}} \mathbf{E}_{\hat{\mathbf{z}}}, \tag{11}$$

where

$$\mathbf{M}_{\mathbf{m},\,\ell} = \frac{\alpha_{\mathbf{m}} \beta_{\,\ell} + k_{\,\ell} k_{\,\mathbf{m}}}{\alpha_{\,\ell} + \beta_{\,\mathbf{m}}} \, (\mathbf{i})^{\mathbf{m} - \ell} \mathbf{J}_{\mathbf{m} - \ell} (\xi \{\alpha_{\,\ell} - \beta_{\,\mathbf{m}}\}) \pi \tag{12}$$

$$\mu_{m} = \frac{-\alpha_{0}\beta_{m} + k_{0}k_{m}}{\alpha_{0} + \beta_{m}} (i)^{m} J(\xi[\alpha_{0} + \beta_{m}]), \qquad (13)$$

and $J_m(x)$ is the m-th-order Bessel function. In general, this results in an infinite number of coupled equations, from which the A_2 's can be extracted. In practice, Agassi and George have shown that 50 equations is sufficient to find an accurate solution. However, in our circumstance this is impossible. We must find the plasmon field strength as a function of both grating height and wavenumber, which means that we would have to invert two 50 x 50 real matrices for each data point. Using the approximation scheme which we shall derive, we have calculated 10,000 data points. This is perhaps more than necessary, but doing so has enabled us to thoroughly understand the behavior of these functions.

For our approximation, we borrow a page from Maradudin⁵ and extend that result to higher orders. The method rests on two ideas. One is that if \mathbf{A}_1 is a plasmon at resonance with \mathbf{k}_g , then

$$A_1 >> A_m \qquad m \neq 1. \tag{14}$$

The second condition is to note that

$$M_{nn} \sim 1$$
, $M_{n,n\pm 1} \sim \xi$, $M_{n,n\pm 2} \sim \xi^2$, ... (15)

so that even for reasonably deep gratings we can assume that ${\rm M}_{11} > {\rm M}_{12} > {\rm M}_{13}$ etc.

A special case of eq. 11 is

$$M_{11}A_{1} + \sum_{m \neq 1} M_{1,m}A_{m} = \mu_{1}E_{1}$$
 (16)

or

$$A_{1} = \frac{\mu_{1} E_{i} + \sum_{m \neq 1} H_{1m} A_{m}}{H_{11}}.$$
 (17)

Thus $R = M_{11}$, which must therefore go to zero if A_1 is to be resonant. Expanding A_{11} to first order in ξ yields our result of eq. 4.

Equation 17 is exact. Now we make our approximation. Suppose that $\mathbf{m} \neq \mathbf{1}$. Then we can write

$$\mathbf{M}_{\mathbf{mm}}\mathbf{A}_{\mathbf{m}} + \sum_{\mathbf{p} \neq \mathbf{m}} \mathbf{M}_{\mathbf{mp}}\mathbf{A}_{\mathbf{p}} = \mu_{\mathbf{m}}\mathbf{E}_{\mathbf{i}}$$
 (18)

But because we are at resonance, of the sum over p, p=1 is by far the largest term. Thus we can simplify eq. 18 as

$$A_{m} = \frac{\mu_{m} E_{1} - M_{m, 1} A_{1}}{M_{mm}}.$$
 (19)

Substituting eq. 19 into 17, we obtain

$$A_{1} = \frac{(\mu_{1} - \sum_{p \neq 1} \frac{M_{1p}}{M_{pp}} \mu_{p}) E_{i}}{M_{11} - \sum_{p \neq 1} \frac{M_{1p}M_{p1}}{M_{pp}}}.$$
 (20)

In the case of normal incidence, $A_1 = A_{-1}$ by symmetry, and we are not justified in leaving A_{-1} out of eq. 19. Inclusion of this term for normal incidence yields

$$A_{1} = \frac{(\mu_{1} - \sum_{p \neq 1} (M_{1p} + M_{1-p}) \frac{\mu_{p}}{M_{pp}}) E_{1}}{[(M_{11} + M_{1-1}) - \sum_{p \neq 1} (M_{1p} + M_{1-p}) (M_{p1} + M_{p-1}) \frac{1}{M_{pp}}}$$
(21)

where p is summed only over non-negative integers. It is clear that in the shallow grating case, both eqs. 20 and 21 reduce to eq. 7.

We shall use eq. 21 with one modification: To ensure numerical convergence, we shall use the dressed Rayleigh expansion. Thus we rewrite eq. 10 as

$$\sum_{\ell=-\infty}^{\infty} M_{m,\ell}^{D} A_{\ell}^{D} = \mu_{m}^{D} E_{i}$$
(22)

where

SERVICE CONTROL OF THE PROPERTY OF THE PROPERT

$$H_{m,n}^{D} = H_{m,n} \exp(i\xi[\alpha_{n} + \beta_{m}])$$
(23)

$$\mu_{\mathbf{m}}^{\mathbf{D}} = \mu_{\mathbf{m}} \exp(\mathrm{i}\beta_{\mathbf{m}}\xi)$$
 , $\mathbf{A}_{\mathbf{n}}^{\mathbf{D}} = \mathbf{A}_{\mathbf{n}} \exp(\mathrm{i}\alpha_{\mathbf{n}}\xi)$.

This eliminates numerical instabilities, not only in the exact calculation, but also for our approximation. In practice, for the grating heights of interest, we have found that summing p from 0 to 15 in eq. 21 provides sufficient accuracy.

APAN MOGERI 1656556 Netroped Telephon Parades (Celebra) Celebras Celebras Celebras Celebras Celebras Celebras

4. Conclusion

Equation 21, modified if necessary by the transformation of eq. 23, constitutes our closed form approximation for the plasmon field strength. Let us consider the advantages and shortcomings of our approximation. The first constraint is to notice that for our approximation to be valid, eq. 14 must hold. This means that we must be on the resonance frequency. Were we not on the resonance frequency, then the approximation used in deriving eq. 19 would not be valid. Secondly, our approximation still is a function of grating height. Note that we have used eq. 15, which means that we are discarding higher orders of ξ for non-resonant plasmons. As these become large, then our approximation becomes less accurate.

Nevertheless, we are happy to report that this method preserves the qualitative features of the full calculation. Figure 1 illustrates this very clearly. Here we have calculated the plasmon field strength ($|A_1|$) for various grating heights using both our new method and the full calculation. It is clear that for gratings up to $\xi \approx 25$ nm, the simple calculation preserves the qualitative features of the exact method. This corresponds to a value of $k_g \xi \approx .7$, which is a very deep grating. The quantitative picture may also be better than appears in the graph. The calculation was done for $k_g \approx 2.95 \times 10^7 \, \mathrm{m}^{-1}$. This is the resonance frequency for shallow gratings, but it is off resonance for deeper gratings. At 25 nm, we have observed that the resonance frequency is closer to 3.1 x 10^{-7} m⁻¹. We have ignored this shift in frequency in our calculations.

As mentioned previously, our result compares with the more general result of Weber. 12 His is more general in that it explicitly accounts for three terms in the Rayleigh expansion rather than just one. However, at resonance, the other two terms can be expected to contribute negligibly, and

hence our result should be nearly as accurate. Off resonance, of course, the other terms are significant and Weber's method is clearly preferable. Strict numerical comparison is impossible since he did his calculation using silver, with a small ε_2 , whereas ours is for cadmium. Furthermore, his example is for a sawtooth grating while ours is for a sinusoidal grating. But it is very clear from a comparision of figure 1 with his table I that the same qualitative features hold, namely that the plasmon resonance intensity is underestimated. Our formalism, while not as general, is derived much more simply and is easier to use, and is probably just as accurate where applicable.

In a later paper 15 we shall apply this formalism to the problem of periodic, laser-induced, chemical vapor deposition. We should be able to predict the growth rate and maximum peak height using the above method.

Acknowledgements

KANM ANDONE AND STREET LAGGEGES BEDEETEN BEGEGESTE BANDON WORKS OF

D. A. J. gratefully acknowledges the invaluable assistance of Dr. Pui-Tak Leung and also appreciates the help of other members of our research group. We would like to thank a referee for bringing refs. 11 and 12 to our attention. This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copywrite notation hereon.


References

- 1. D.J. Ehrlich and J.Y. Tsao, J. Vac. Sci. Technol. B 1, 4, 969 (1983), and references therein.
- 2. R.M. Osgood, Jr., and D.J. Ehrlich, Opt. Lett. 7, 385 (1982).
- 3. S.R.J. Brueck and D.J. Ehrlich, Phys. Rev. Lett. 48, 1678 (1982).
- 4. R. Petit, in <u>Electromagnetic Theory of Gratings</u>, edited by R. Petit (Springer-Verlag, Berlin, 1980), Chapt. 1.
- 5. A.A. Maradudin, in <u>Surface Polaritons</u>, edited by V.M. Agranovich and D.L. Mills (North-Holland, Amsterdam, 1982), Chapt. 10.
- 6. S.S. Jha, J.R. Kirtley and J.C. Tsang, Phys. Rev. B <u>22</u>, 3973 (1980).
- F. Toigo, A. Marvin, V. Celli and N.R. Hill, Phys. Rev. B <u>15</u>, 5618 (1977).
- 8. A. Marvin, F. Toigo and V. Celli, Phys. Rev. B <u>11</u>, 2777 (1975).
- 9. D. Agassi and T.F. George, Phys. Rev. B 33, 2393 (1986).
- 10. D. Agassi and T.F. George, Surf. Sci. 172, 230 (1986).
- 11. N.E. Glass, M. Weber and D.L. Mills, Phys. Rev. B 29, 6548 (1984).
- 12. M.G. Weber, Phys. Rev. B <u>33</u>, 9, (1986).
- P.M. Fauchet and A.E. Siegman, Appl. Phys. Lett. 40, 824 (1982).
- J.E. Sipe, J.F. Young, J.S. Preston and H.M. van Driei, Phys. Rev. B 27, 1141 (1983).
- 15. D.A. Jelski and T.F. George, J. Appl. Phys., in press.

FIGURE CAPTION

Figure 1. This graph compares the plasmon field strength ($|A_1|$) calculated from eq. 21 (using the dressed Rayleigh expansion) with that from the exact method. The dielectric constant used is -2.5 + 1.3i, the incident light has a wavelength of 257 nm ($k = 2.44 \times 10^7 \text{ m}^{-1}$), and the grating wavenumber is taken as 2.95 $\times 10^7 \text{ m}^{-1}$.

SSSS SOCIOCO COSSSS SCOREGO SSSSSSSS CONTRACTOR CONTRAC



01/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Washington, D.C. 20375-5000	. •	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis. Minnesota 55455

Dr. A. Reisman Microelectronics Center of North Carolina Research Triangle Park, North Carolina 27709

Dr. M. Grunze Laboratory for Surface Science and Technology University of Maine Orono, Maine 04469

Dr. J. Butler Naval Research Laboratory Code 6115 Washington D.C. 20375-5000

Dr. L. Interante Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Irvin Heard Chemistry and Physics Department Lincoln University Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde Department of Chemistry Kansas State University Manhattan, Kansas 66506 Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookesville, Tennessee 38501

Dr. D. Dileila Chemistry Department George Washington University Washington D.C. 20052

Dr. R. Reeves Chemistry Department Renssaeler Polytechnic Institute Troy, New York 12181

Dr. Steven M. George Stanford University Department of Chemistry Stanford, CA 94305

Dr. Mark Johnson Yale University Department of Chemistry New Haven, CT 06511-8118

Dr. W. Knauer Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Cr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60637

Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712

Dr. D. E. Harrison Department of Physics Naval Postgraduate School Monterey, California 93940 Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green Quantum Surface Dynamics Branch Code 3817 Naval Weapons Center China Lake, California 93555

Dr. A. Wold Department of Chemistry Brown University Providence, Rhode Island 02912

Dr. S. L. Bernasek Department of Chemistry Princeton University Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Richard Colton Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Dan Pierce National Bureau of Standards Optical Physics Division Washington, D.C. 20234

Dr. R. Stanley Williams Department of Chemistry University of California Los Angeles, California 90024

Dr. R. P. Messmer Materials Characterization Lab. General Electric Company Schenectady, New York 22217

Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637

Dr. Ronald Lee R301 Naval Surface Weapons Center White Oak Silver Spring, Maryland 20910

Dr. Paul Schoen Code 6190 Waval Research Laboratory Washington, D.C. 20375-5000 Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260

Dr. Richard Greene Code 5230 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda University of Pittsburg Chemistry Building Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH UNITED KINGDOM

Dr. H. Tachikawa Chemistry Department Jackson State University Jackson, Mississippi 39217

Dr. John W. Wilkins Cornell University Laboratory of Atomic and Solid State Physics Ithaca, New York 14853

KANARA PYYYYY

TANKS TO SEE THE PARTY OF THE P

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis Department of Physics University of California Irvine, California 92664

Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052

Dr. J. C. Hemminger . Chemistry Department University of California Irvine, California 92717

Dr. T. F. George Chemistry Department University of Rochester Rochester, New York 14627

Dr. G. Rubloff IBM Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. Horia Metiu Chemistry Department University of California Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma Department of Physics University of California Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser Department of Chemistry University of Richmond Richmond, Virginia 23173

Dr. R. W. Plummer Department of Physics University of Pennsylvania Philadelphia, Pennsylvania 19104

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann Department of Chemistry Cornell University Ithaca, New York 14853

Or. A. Steck1
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, NewYork 12181

Dr. G.H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853